











Rules for Derivatives and Other Useful Facts

$(x^n)' = nx^{n-1}$ for any real n	$(\sqrt{x})' = \frac{1}{2\sqrt{x}}$	$(\sin x)' = \cos x$ $(\cos x)' = -\sin x$
$(b^x)' = (\ln b)b^x$ for any $b > 0$	$(e^x)' = e^x$	$(\ln x)' = \frac{1}{x}$
$(cf)' = cf'$ for constant c $(f \pm g)' = f' \pm g'$	$[f(g(x))]' = f'(g(x))g'(x)$	$(fg)' = f'g + g'f$ $\left(\frac{f}{g}\right)' = \frac{f'g - g'f}{g^2}$

Trigonometric Identities

$\tan x = \frac{\sin x}{\cos x}$	$\cot x = \frac{\cos x}{\sin x}$	$\sec x = \frac{1}{\cos x}$	$\csc x = \frac{1}{\sin x}$	$\sin^2 x + \cos^2 x = 1$
----------------------------------	----------------------------------	-----------------------------	-----------------------------	---------------------------

Geometry: Angles in Radians; P = perimeter, A = area or surface area, V = volume

<p>TRIANGLE</p> <p>$P = a + b + c$ $A = \frac{1}{2}bh$</p> 	<p>RECTANGLE</p> <p>$P = 2a + 2b$ $A = ab$</p> 	<p>CIRCLE</p> <p>$P = 2\pi r$ $A = \pi r^2$</p> 
<p>PYTHAGOREAN THEOREM</p> <p>$a^2 + b^2 = c^2$ $c = \sqrt{a^2 + b^2}$</p> 	<p>TRAPEZOID</p> <p>$P = a + b + c + d$ $A = h \frac{a+b}{2}$</p> 	<p>CIRCULAR SECTOR</p> <p>$L = r\theta$ $A = \frac{1}{2}r^2\theta$</p> 
<p>RECTANGULAR BOX</p> <p>$A = 2ab + 2ac + 2bc$ $V = abc$</p> 	<p>SQUARE PYRAMID</p> <p>$A = s(s + \sqrt{s^2 + 4h^2})$ $V = \frac{1}{3}s^2h$</p> 	<p>RIGHT CIRCULAR CONE</p> <p>$A = \pi r^2 + \pi rs$ $s = \sqrt{r^2 + h^2}$ $V = \frac{1}{3}\pi r^2 h$</p> 
<p>CYLINDER</p> <p>$A = 2\pi r(r + h)$ $V = \pi r^2 h$</p> 	<p>SPHERE</p> <p>$A = 4\pi r^2$ $V = \frac{4\pi r^3}{3}$</p> 